



Nadolol


Dosage Form: tablet

Nadolol TABLETS USP

Rx only

Nadolol Description

Nadolol USP is a synthetic nonselective beta-adrenergic receptor blocking agent designated chemically as 1-(tert-butylamino)-3-[(5,6,7,8-tetrahydro-cis-6,7-dihydroxy-1-naphthyl)oxy]-2-propanol. The structural formula is:



C₁₇H₂₇NO₄

M.W. 309.40

Nadolol USP is a white to off-white, practically odorless, crystalline powder. It is freely soluble in water at pH2, in alcohol, and in methanol and slightly soluble in chloroform. Nadolol USP is available for oral administration as 20 mg, 40 mg, and 80 mg tablets and contains the following inactive ingredients: citric acid, corn starch, magnesium stearate, microcrystalline cellulose and povidone. In addition, the 80 mg tablets contain sodium starch glycolate.

Nadolol - Clinical Pharmacology

Nadolol is a nonselective beta-adrenergic receptor blocking agent. Clinical pharmacology studies have demonstrated beta-blocking activity by showing (1) reduction in heart rate and cardiac output at rest and on exercise, (2) reduction of systolic and diastolic blood pressure at rest and on exercise, (3) inhibition of isoproterenol-induced tachycardia, and (4) reduction of reflex orthostatic tachycardia.

Nadolol specifically competes with beta-adrenergic receptor agonists receptors located chiefly in cardiac muscle and the beta2 receptors located chiefly in the bronchial and vascular musculature, inhibiting the chronotropic, inotropic, and vasodilator responses to beta-adrenergic stimulation proportionately. Nadolol has no intrinsic sympathomimetic activity and, unlike some other beta-adrenergic blocking agents, Nadolol has little direct myocardial depressant activity and does not have an anesthetic-like membrane-stabilizing action. Animal and human studies show that Nadolol slows the sinus rate and depresses AV conduction. In dogs, only minimal amounts of Nadolol were detected in the brain relative to amounts in blood and other organs and tissues. Nadolol has low lipophilicity as determined by octanol/water partition coefficient, a characteristic of certain beta-blocking agents that has been correlated with the limited extent to which these agents cross the blood-brain barrier, their low concentration in the brain, and low incidence of CNS-related side effects.

In controlled clinical studies, Nadolol at doses of 40 to 320 mg/day has been shown to decrease both standing and supine blood pressure, the effect persisting for approximately 24 hours after dosing.

The mechanism of the antihypertensive effects of beta-adrenergic receptor blocking agents has not been established; however, factors that may be involved include (1) competitive antagonism of catecholamines at peripheral (non-CNS) adrenergic neuron sites (especially cardiac) leading to decreased cardiac output, (2) a central effect leading to reduced tonic-sympathetic nerve outflow to the periphery, and (3) suppression of renin secretion by blockade of the beta-adrenergic receptors responsible for renin release from the kidneys.

While cardiac output and arterial pressure are reduced by Nadolol therapy, renal hemodynamics are stable, with preservation of renal blood flow and glomerular filtration rate.

By blocking catecholamine-induced increases in heart rate, velocity and extent of myocardial contraction, and blood pressure, Nadolol generally reduces the oxygen requirements of the heart at any given level of effort, making it useful for many patients in the long-term management of angina pectoris. On the other hand, Nadolol can increase oxygen requirements by increasing left ventricular fiber length and end diastolic pressure, particularly in patients with heart failure.

Although beta-adrenergic receptor blockade is useful in treatment of angina and hypertension, there are also situations in which sympathetic stimulation is vital. For example, in patients with severely damaged hearts, adequate ventricular function may depend on sympathetic drive. Beta-adrenergic blockade may worsen AV block by preventing the necessary facilitating effects of sympathetic activity on conduction. Beta2-adrenergic blockade results in passive bronchial constriction by interfering with endogenous adrenergic bronchodilator activity in patients subject to bronchospasm and may also interfere with exogenous bronchodilators in such patients.

Absorption of Nadolol after oral dosing is variable, averaging about 30 percent. Peak serum concentrations of Nadolol usually occur in three to four hours after oral administration and the presence of food in the gastrointestinal tract does not affect the rate or extent of Nadolol absorption. Approximately 30 percent of the Nadolol present in serum is reversibly bound to plasma protein.

Unlike many other beta-adrenergic blocking agents, Nadolol is not metabolized by the liver and is excreted unchanged, principally by the kidneys.

The half-life of therapeutic doses of Nadolol is about 20 to 24 hours, permitting once-daily dosage. Because Nadolol is excreted predominantly in the urine, its half-life increases in renal failure (see PRECAUTIONS and DOSAGE AND ADMINISTRATION). Steady-state serum concentrations of Nadolol are attained in six to nine days with once-daily dosage in persons with normal renal function. Because of variable absorption and different individual responsiveness, the proper dosage must be determined by titration.

Exacerbation of angina and, in some cases, myocardial infarction and ventricular dysrhythmias have been reported after abrupt discontinuation of therapy with beta-adrenergic blocking agents in patients with coronary artery disease. Abrupt withdrawal of these agents in patients without coronary artery disease has resulted in transient symptoms, including tremulousness, sweating, palpitation, headache, and malaise. Several mechanisms have been proposed to explain these phenomena, among them increased sensitivity to catecholamines because of increased numbers of beta receptors.

Indications and Usage for Nadolol

Did you find this page helpful? Yes No

Angina Pectoris

Nadolol tablets are indicated for the long-term management of patients with angina pectoris.

Hypertension

Nadolol tablets are indicated in the management of hypertension; they may be used alone or in combination with other antihypertensive agents, especially thiazide-type diuretics.

Contraindications

Nadolol is contraindicated in bronchial asthma, sinus bradycardia and greater than first degree conduction block, cardiogenic shock, and overt cardiac failure (see WARNINGS).

Warnings

Cardiac Failure

Sympathetic stimulation may be a vital component supporting circulatory function in patients with congestive heart failure, and its inhibition by beta-blockade may precipitate more severe failure. Although beta-blockers should be avoided in overt congestive heart failure, if necessary, they can be used with caution in patients with a history of failure who are well-compensated, usually with digitalis and diuretics. Beta-adrenergic blocking agents do not abolish the inotropic action of digitalis on heart muscle.

IN PATIENTS WITHOUT A HISTORY OF HEART FAILURE, continued use of beta-blockers can, in some cases, lead to cardiac failure. Therefore, at the first sign or symptom of heart failure, the patient should be digitalized and/or treated with diuretics, and the response observed closely, or Nadolol should be discontinued (gradually, if possible).

Exacerbation of Ischemic Heart Disease Following Abrupt Withdrawal

Hypersensitivity to catecholamines has been observed in patients withdrawn from beta-blocker therapy; exacerbation of angina and, in some cases, myocardial infarction have occurred after abrupt discontinuation of such therapy. When discontinuing chronically administered Nadolol, particularly in patients with ischemic heart disease, the dosage should be gradually reduced over a period of one to two weeks and the patient should be carefully monitored. If angina markedly worsens or acute coronary insufficiency develops, Nadolol administration should be reinstated promptly, at least temporarily, and other measures appropriate for the management of unstable angina should be taken. Patients should be warned against interruption or discontinuation of therapy without the physician's advice. Because coronary artery disease is common and may be unrecognized, it may be prudent not to discontinue Nadolol therapy abruptly even in patients treated only for hypertension.

Nonallergic Bronchospasm (e.g., chronic bronchitis, emphysema)

PATIENTS WITH BRONCHOSPASTIC DISEASES SHOULD IN GENERAL NOT RECEIVE BETA-BLOCKERS. Nadolol should be administered with caution since it may block bronchodilation produced by endogenous or exogenous catecholamine stimulation of beta₂ receptors.

Major Surgery

Chronically administered beta-blocking therapy should not be routinely withdrawn prior to major surgery; however, the impaired ability of the heart to respond to reflex adrenergic stimuli may augment the risks of general anesthesia and surgical procedures.

Diabetes and Hypoglycemia

Beta-adrenergic blockade may prevent the appearance of premonitory signs and symptoms (e.g., tachycardia and blood

pressure changes) of acute hypoglycemia. This is especially important with labile diabetics. Beta-blockade also reduces the release of insulin in response to hyperglycemia; therefore, it may be necessary to adjust the dose of antidiabetic drugs.

Thyrotoxicosis

Beta-adrenergic blockade may mask certain clinical signs (e.g., tachycardia) of hyperthyroidism. Patients suspected of developing thyrotoxicosis should be managed carefully to avoid abrupt withdrawal of beta-adrenergic blockade which might precipitate a thyroid storm.

Precautions

Impaired Renal Function

Nadolol should be used with caution in patients with impaired renal function (see DOSAGE AND ADMINISTRATION).

Information for Patients

Patients, especially those with evidence of coronary artery insufficiency, should be warned against interruption or discontinuation of Nadolol therapy without the physician's advice. Although cardiac failure rarely occurs in properly selected patients, patients being treated with beta-adrenergic blocking agents should be advised to consult the physician at the first sign or symptom of impending failure. The patient should also be advised of a proper course in the event of an inadvertently missed dose.

Drug Interactions

When administered concurrently, the following drugs may interact with beta-adrenergic receptor blocking agents:

Anesthetics, General

Exaggeration of the hypotension induced by general anesthetics (see WARNINGS, Major Surgery).

Antidiabetic Drugs (oral agents and insulin)

Hypoglycemia or hyperglycemia; adjust dosage of antidiabetic drug accordingly (see WARNINGS, Diabetes and Hypoglycemia).

Catecholamine-depleting Drugs (e.g., reserpine)

Additive effect; monitor closely for evidence of hypotension and/or excessive bradycardia (e.g., vertigo, syncope, postural hypotension).

Digitalis Glycosides

Both digitalis glycosides and beta-blockers slow atrioventricular conduction and decrease heart rate. Concomitant use can increase the risk of bradycardia.

Response to Treatment for Anaphylactic Reaction

While taking beta-blockers, patients with a history of severe anaphylactic reaction to a variety of allergens may be more reactive to repeated challenge, either accidental, diagnostic, or therapeutic. Such patients may be unresponsive to the usual doses of epinephrine used to treat allergic reaction.

Carcinogenesis, Mutagenesis, Impairment of Fertility

In chronic oral toxicologic studies (one to two years) in mice, rats, and dogs, Nadolol did not produce any significant toxic effects. In two year oral carcinogenic studies in rats and mice, Nadolol did not produce any neoplastic, preneoplastic, or non-neoplastic pathologic lesions. In fertility and general reproductive performance studies in rats, Nadolol caused no adverse effects.

Pregnancy

Teratogenic Effects

Pregnancy Category C

In animal reproduction studies with Nadolol, evidence of embryo- and fetotoxicity was found in rabbits, but not in rats or hamsters, at doses 5 to 10 times greater (on a mg/kg basis) than the maximum indicated human dose. No teratogenic potential was observed in any of these species. There are no adequate and well-controlled studies in pregnant women. Nadolol should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Neonates whose mothers are receiving Nadolol at parturition have exhibited bradycardia, hypoglycemia, and associated symptoms.

Nursing Mothers

Nadolol is excreted in human milk. Because of the potential for adverse effects in nursing infants, a decision should be made whether to discontinue nursing or to discontinue therapy taking into account the importance of Nadolol to the mother.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

Adverse Reactions

Most adverse effects have been mild and transient and have rarely required withdrawal of therapy.

Cardiovascular

Bradycardia with heart rates of less than 60 beats per minute occurs commonly, and heart rates below 40 beats per minute and/or symptomatic bradycardia were seen in about 2 of 100 patients. Symptoms of peripheral vascular insufficiency, usually of the Raynaud type, have occurred in approximately 2 of 100 patients. Cardiac failure, hypotension, and rhythm/conduction disturbances have each occurred in about 1 of 100 patients. Single instances of first degree and third degree heart block have been reported; intensification of AV block is a known effect of beta-blockers (see also CONTRAINDICATIONS, WARNINGS, and PRECAUTIONS).

Central Nervous System

Dizziness or fatigue has been reported in approximately 2 of 100 patients; paresthesias, sedation, and change in behavior have each been reported in approximately 6 of 1000 patients.

Respiratory

Bronchospasm has been reported in approximately 1 of 1000 patients (see CONTRAINDICATIONS and WARNINGS).

Gastrointestinal

Nausea, diarrhea, abdominal discomfort, constipation, vomiting, indigestion, anorexia, bloating, and flatulence have been reported in 1 to 5 of 1000 patients.

Miscellaneous

Each of the following has been reported in 1 to 5 of 1000 patients: rash; pruritus; headache; dry mouth, eyes, or skin; impotence or decreased libido; facial swelling; weight gain; slurred speech; cough; nasal stuffiness; sweating; tinnitus; blurred vision. Reversible alopecia has been reported infrequently.

The following adverse reactions have been reported in patients taking Nadolol and/or other beta-adrenergic blocking agents, but no causal relationship to Nadolol has been established.

Central Nervous System

Reversible mental depression progressing to catatonia; visual disturbances; hallucinations; an acute reversible syndrome characterized by disorientation for time and place, short-term memory loss, emotional lability with slightly clouded sensorium, and decreased performance on neuropsychometrics.

Gastrointestinal

Mesenteric arterial thrombosis; ischemic colitis; elevated liver enzymes.

Hematologic

Agranulocytosis; thrombocytopenic or nonthrombocytopenic purpura.

Allergic

Fever combined with aching and sore throat; laryngospasm; respiratory distress.

Miscellaneous

Pemphigoid rash; hypertensive reaction in patients with pheochromocytoma; sleep disturbances; Peyronie's disease.

The oculomucocutaneous syndrome associated with the beta-blocker practolol has not been reported with Nadolol.

Overdosage

Nadolol can be removed from the general circulation by hemodialysis.

In addition to gastric lavage, the following measures should be employed, as appropriate. In determining the duration of corrective therapy, note must be taken of the long duration of the effect of Nadolol.

Excessive Bradycardia

Administer atropine (0.25 to 1 mg). If there is no response to vagal blockade, administer isoproterenol cautiously.

Cardiac Failure

Administer a digitalis glycoside and diuretic. It has been reported that glucagon may also be useful in this situation.

Hypotension

Administer vasopressors, e.g., epinephrine or levarterenol. (There is evidence that epinephrine may be the drug of choice.)

Bronchospasm

Administer a beta2-stimulating agent and/or a theophylline derivative.

Nadolol Dosage and Administration

DOSAGE MUST BE INDIVIDUALIZED. Nadolol MAY BE ADMINISTERED WITHOUT REGARD TO MEALS.

Angina Pectoris

The usual initial dose is 40 mg Nadolol once daily. Dosage may be gradually increased in 40 to 80 mg increments at 3 to 7 day intervals until optimum clinical response is obtained or there is pronounced slowing of the heart rate. The usual maintenance dose is 40 or 80 mg administered once daily. Doses up to 160 or 240 mg administered once daily may be needed.

The usefulness and safety in angina pectoris of dosage exceeding 240 mg per day have not been established. If treatment is to be discontinued, reduce the dosage gradually over a period of one to two weeks (see WARNINGS).

Hypertension

The usual initial dose is 40 mg Nadolol once daily, whether it is used alone or in addition to diuretic therapy. Dosage may be gradually increased in 40 to 80 mg increments until optimum blood pressure reduction is achieved. The usual maintenance dose is 40 or 80 mg administered once daily. Doses up to 240 or 320 mg administered once daily may be needed.

Dosage Adjustment in Renal Failure

Absorbed Nadolol is excreted principally by the kidneys and, although nonrenal elimination does occur, dosage adjustments are necessary in patients with renal impairment. The following dose intervals are recommended:

Creatinine Clearance (mL/min/1.73m ²)	Dosage Interval (hours)
> 50	24
31 to 50	24 to 36
10 to 30	24 to 48
< 10	40 to 60

How is Nadolol Supplied

Nadolol Tablets USP, 20 mg are available as white, round tablet debossed "20" on one side, and a bisect on the other side with "Z" on the upper half and "4235" on the lower half, containing 20 mg of Nadolol USP packaged in bottles of 100 tablets.

Nadolol Tablets USP, 40 mg are available as white, round tablet debossed "40" on one side, and a bisect on the other side with "Z" on the upper half and "4236" on the lower half, containing 40 mg of Nadolol USP packaged in bottles of 100 tablets.

Nadolol Tablets USP, 80 mg are available as white, round tablet debossed "80" on one side, and a bisect on the other side with "Z" on the upper half and "4237" on the lower half, containing 80 mg of Nadolol USP packaged in bottles of 100 tablets.

Dispense in a tight container as defined in the USP, with a child-resistant closure (as required).

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Manufactured In India By:

EMCURE PHARMACEUTICALS LTD.

Hinjwadi, Pune, India

Manufactured For:

TEVA PHARMACEUTICALS USA

Sellersville, PA 18960

Iss. 6/2011

PRINCIPAL DISPLAY PANEL



Each tablet contains:
nadolol, USP 20 mg

Usual Dosage: See package insert for full prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a tight container as defined in the USP, with a child-resistant closure (as required).

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Code No.: MH/DRUGS/PD/149 Iss. 7/2007

Manufactured In India By:
EMCURE PHARMACEUTICALS LTD.
Hinjwadi, Pune, India

Manufactured For:
TEVA PHARMACEUTICALS USA
Sellersville, PA 18960



Nadolol Tablets USP 20 mg 100s Label Text

NDC 0093-4235-01

Nadolol
 Tablets USP
 20 mg
 Rx only
 100 TABLETS
 TEVA

PRINCIPAL DISPLAY PANEL

NDC 0093-4236-01
NADOLOL
Tablets USP
40 mg

Each tablet contains:
 nadolol, USP

Rx only

100 TABLETS

TEVA

40 mg

LOT:
EXP:

0093-4236-01

5102700221001

Usual Dosage: See package insert for full prescribing information.
 Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].
 Dispense in a tight container as defined in the USP, with a child-resistant closure (as required).
KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.
 Code No.: MH/DRUGS/PD/149 Iss. 7/2007
 Manufactured in India By:
 EMCURE PHARMACEUTICALS LTD.
 Hinjwadi, Pune, India
 Manufactured For:
TEVA PHARMACEUTICALS USA
 Sellersville, PA 18960

Nadolol Tablets USP 40 mg 100s Label Text

NDC-0093-4236-01
 Nadolol
 Tablets USP
 40 mg
 Each tablet contains:
 Nadolol, USP 40 mg
 Rx only
 100 TABLETS
 TEVA

PRINCIPAL DISPLAY PANEL

NDC 0093-4237-01
NADOLOL
Tablets USP
80 mg

Each tablet contains:
 nadolol, USP

Rx only

100 TABLETS

TEVA

80 mg

LOT:
EXP:

0093-4237-01

5102700231001

Usual Dosage: See package insert for full prescribing information.
 Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].
 Dispense in a tight container as defined in the USP, with a child-resistant closure (as required).
KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.
 Code No.: MH/DRUGS/PD/149 Iss. 7/2007
 Manufactured in India By:
 EMCURE PHARMACEUTICALS LTD.
 Hinjwadi, Pune, India
 Manufactured For:
TEVA PHARMACEUTICALS USA
 Sellersville, PA 18960

Nadolol Tablets USP 80 mg 100s Label Text

NDC 0093-4237-01

Nadolol

Tablets USP

80 mg

Each tablet contains:

Nadolol, USP 80 mg

Rx only

100 TABLETS

TEVA

**Nadolol
Nadolol tablet****Product Information****Product Type** HUMAN PRESCRIPTION DRUG **NDC Product Code (Source)** 0093-4235**Route of Administration** ORAL **DEA Schedule****Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
Nadolol (Nadolol)	Nadolol	20 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE	
STARCH, CORN	
MAGNESIUM STEARATE	
CELLULOSE, MICROCRYSTALLINE	
POVIDONE	

Product Characteristics**Color** WHITE **Score** 2 pieces

Shape	ROUND	Size	8mm
Flavor		Imprint Code	20;Z;4235
Contains			

Packaging

#	NDC	Package Description	Multilevel Packaging
1	0093-4235-01	100 TABLET In 1 BOTTLE	None

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074229	01/18/2011	

**Nadolol
Nadolol tablet****Product Information**

Product Type	HUMAN PRESCRIPTION DRUG	NDC Product Code (Source)	0093-4236
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Nadolol (Nadolol)	Nadolol	40 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE	
STARCH, CORN	
MAGNESIUM STEARATE	
CELLULOSE, MICROCRYSTALLINE	
POVIDONE	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	40;Z;4236

Contains**Packaging**

#	NDC	Package Description	Multilevel Packaging
1	0093-4236-01	100 TABLET In 1 BOTTLE	None

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074229	01/18/2011	

**Nadolol
Nadolol tablet****Product Information**

Product Type	HUMAN PRESCRIPTION DRUG	NDC Product Code (Source)	0093-4237
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Nadolol (Nadolol)	Nadolol	80 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE	
STARCH, CORN	
MAGNESIUM STEARATE	

CELLULOSE, MICROCRYSTALLINE

POVIDONE

SODIUM STARCH GLYCOLATE TYPE A POTATO

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	11mm
Flavor		Imprint Code	80;Z;4237

Contains

Packaging

#	NDC	Package Description	Multilevel Packaging
1	0093-4237-01	100 TABLET In 1 BOTTLE	None

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074255	01/18/2011	

Labeler - Teva Pharmaceuticals USA Inc (118234421)

Revised: 01/2011 Teva Pharmaceuticals USA Inc